


Original Research Article

Evaluation of breast lesions by fine needle aspiration cytology

Ayesha aka Arnaz Khan^{1*}, Kevan Kothari², S.S. Goswami³

^{1,2,3}Smt. B.K. Shah Medical Institute and Research Centre, Piparia, Vadodara, Gujarat, India

*Corresponding author email: 1993arnaz@gmail.com

	International Archives of Integrated Medicine, Vol. 8, Issue 12, December, 2021. Available online at http://iaimjournal.com/ ISSN: 2394-0026 (P) ISSN: 2394-0034 (O)
	Received on: 02-12-2021 Accepted on: 11-12-2021 Source of support: Nil Conflict of interest: None declared. Article is under creative common license CC-BY
How to cite this article: Ayesha aka Arnaz Khan, Kevan Kothari, S.S. Goswami. Evaluation of breast lesions by fine needle aspiration cytology. IAIM, 2021; 8(12): 68-73.	

Abstract

Background: FNAC is very useful technique and routinely done on palpable lesions of the body as a diagnostic procedure. For preoperative diagnosis of breast lesions, triple approach technique is used in which FNAC is one of the most important techniques. The main purpose of FNAC of breast lesions is in the investigation of any palpable lump and to avoid unnecessary surgery in specific benign conditions. The advantages are- it provides rapid and accurate diagnosis, is therapeutic as well as diagnostic in many cystic conditions.

Materials and methods: The study was community based cross sectional study and was done at Pathology Department of SBKS MI & RC, Vadodara, Gujarat over a period of 18 months from January 2019 to July 2020.

Results: The C2 category of 60 benign cases consisted of 20 (majority) cases of fibroadenoma, 3 of phyllodes tumor, 15 of benign proliferative breast disease, 1 of fibrocystic breast disease, 7 of inflammatory lesions and 3 of intraductal papilloma. Out of 5 cases of gynecomastia, 4 were benign and 1 was malignant (intracystic papillary carcinoma) which confirmed by histopathology. The C5 category of 32 malignant lesions consisted of 31 cases of infiltrating duct carcinoma- NOS type and 1 of infiltrating duct carcinoma with medullary type.

Conclusion: The study was 100% sensitive in the diagnosis of malignant lesions. The higher incidence of malignant breast lesions with aged group 41– 60 years of patients. The C2 category of 60 benign cases consisted of 20 (majority) cases of fibroadenoma. The C5 category of 32 malignant lesions consisted of 31 cases of infiltrating duct carcinoma- NOS type.

Key words

FNAC, Fine Needle Aspiration Cytology, Breast lesions.

Introduction

Fine needle aspiration cytology (FNAC) is very useful technique and routinely done on palpable lesions such as superficial growths of the skin, subcutis, soft tissues and organs such as thyroid, breast, salivary glands, and superficial lymph nodes. It can be used on all organs of the body as a diagnostic procedure.

Breast carcinoma is the most common in women worldwide and it is the most important cause of death related to cancer for women between ages 20 to 59 years.

For preoperative diagnosis of breast lesions, triple approach technique is used and in this FNAC is one of the most important techniques [1].

The main purpose of FNAC of breast lesions is in the investigation of any palpable lump and to avoid unnecessary surgery in specific benign conditions. The advantages are- it provides rapid and accurate diagnosis, is therapeutic as well as diagnostic in many cystic conditions [2].

The limitation of an FNAC is its inability to separate in situ and invasive carcinoma [3].

However, FNAC still happens to be the popular modality of diagnosis because of its overall accuracy in experienced hands, least invasiveness, ability for repeat testing and speed of giving results.

Aim and objective

- To establish utility and effectiveness in diagnosis of breast lesion by FNAC.
- To study cytological findings of benign and malignant features of breast lesions which co-relate with histopathological findings.
- Find the incidence of breast cancer in palpable lump in different age group.
- For utility of FNAC in various breast lesions at different age group.
- To study pre-operative confirmation of

clinically suspected cancer for guide to further clinical management.

Materials and methods

Study design: Community based cross sectional study.

Study site: Dhiraj General Hospital and Shrimati Bhikhiben Kanjibhai Shah Medical Institute and Research Centre, Sumandeep Vidyapeeth, Piparia, Vadodara, Gujarat.

Study population: All the patients having breast lump referred by department of surgery, Dhiraj Hospital, either indoor or outpatient basis.

Sample size: 100 cases were carried out.

Inclusion criteria:

- All the patients referred to cytology section of the Pathology Department, SBKS MI & RC, Piparia, Vadodara, Gujarat.
- Clinically having unilateral or bilateral breast lump.
- Male and Female.
- Patient with breast lesion sent for FNAC and also included USG guided /CT guided FNAC for same.

Exclusion criteria:

- Non cooperative patients.

Results

The present study included FNAC material from the 100 cases of breast lesions (110 aspirates) from January 2019 to July 2020 at the Department of Pathology, SBKS MI & RC. The observations of the study were as follows:

Age and sex distribution

The age of the patients ranged from 11 to 80 years. There were 95 female patients and 5 male patients. The age distribution in relation to sex was shown in **Table - 1**.

Distribution of breast cases in relation to right or left breast

Out of 100 cases of breast lesions, right side involved in 53 cases, left side involved in 41 cases and 6 cases show bilateral involvement.

Table – 1: Age and sex distribution.

Age in year	<=20	21-40	41-60	61-80	Total
Males	1	2	1	1	5
Females	17	46	23	9	95
Total	18	48	24	10	100


Table - 2: Diagnosis of benign breast lesions (C2) with biopsy correlation total 60 cases.

Benign lesions	No of cases	Percentage (benign lesions)	No. of biopsy	Histopathology diagnosis confirmed
Inflammatory lesions	7	11.66%	07	
• Breast abscess	3			Breast abscess (3)
• Acute mastitis	2			Acute mastitis (2)
• Granulomatous mastitis	2			Granulomatous mastitis (2)
Fibroadenoma	20	33.33%	20	Fibroadenoma (20)
Phyllodes tumor	3	5%	03	Phyllodes tumor (3)
Benign proliferative breast disease	21	35%	15	Benign proliferative breast disease (15)
Fibrocystic breast disease	01	1.66%	01	Fibrocystic breast Disease (1)
Intraductal papilloma	03	1.66%	03	Intraductal papilloma (3)
Gynecomastia	05	11.66%	05	Gynecomastia (4)
Total	60		54	

Table - 3: Diagnosis of malignant breast lesions (C5) with biopsy correlation.

Malignant lesions diagnosed at cytology	No. of Cases	Percentage (out of 30 cases)	No. of biopsied	Histopathology diagnosis
Ductal Carcinoma	31	96.87%	31	Infiltrating ductal carcinoma-NOS type
Ductal Carcinoma	1	3.13%	1	Infiltrating ductal carcinoma with Medullary type
Total	32			

Figure - 1: Fibroadenoma (H & E, 4X).



Clinical history

Most of the patients were presented with mass lesions in the breast. 17 cases in addition had

associated pain, 29 cases presented with anorexia, weight loss, fatigue, 3 cases presented with fever, 1 case showed evidence of cracking of nipple underlying to the mass and 1 case show blood mixed nipple discharge.

Adequacy of the sample

Out of the 100 cases of breast lesion, only 1 case was inadequate for aspiration. Of the 99 adequate aspirates, 58 were moderately cellular, and 37 were highly cellular. Only 90 cases underwent a biopsy and could be correlated histopathologically.

Figure - 2: Phyllodes tumor (H&E, 4X): Theand in singly. Cells showed an enlarged increased stromal cellularity.





Figure - 3: Ductal carcinoma (H & E, 40): Malignant cells with clumped chromatin.



NHSBSP (National Health Service of Breast Screening Programme) reporting criteria of FNAC of breast lesions

According to NHSBSP, the cytology of 100 cases were analyzed and categorized into 5 categories from C1 to C5. There was 1 case in C1 category, 60 cases in C2 category, 4 cases in C3 category, 3 case in C4 category and 32 cases in C5 category.

Benign lesions (C2: category)

Out of the 60 benign cases, majority of cases (28 cases) were from upper outer quadrant (**Table - 2, Figure - 1, 2**).

Malignant lesions (C5 category)

Out of the 32 malignant cases, majority of the cases (25 cases) were involved the upper outer quadrant (**Table - 3, Figure - 3**).

Infiltrating duct carcinoma - NOS

Out of the 32 malignant lesions, 31 were reported in cytology as ductal carcinoma. The age ranged from 35 to 80 years. The smears were moderately to highly cellular with pleomorphic malignant ductal epithelial cells in loose clusters, sheets,

hyperchromatic nuclei, increased N:C ratio, moderate to marked nuclear pleomorphism, coarse to granular to clumped chromatin with prominent nucleoli and cytoplasm was eosinophilic. The patients underwent excision of the mass and the histopathology confirmed infiltrating ductal carcinoma (NOS type) in all 31 cases.

Discussion

The age of the patients ranged up to 80 years with the majority in the 21-40 years age group. Out of 100 cases, majority of the breast masses were located in the right breast in the upper outer quadrant and least in the lower outer and inner quadrant.

Out of 100 cases, in 99 cases the aspirates were adequate and 1 was inadequate for interpretation.

In present study, 60 cases were diagnosed benign at cytology. Based on this patients were underwent surgery and histopathology showed 59 cases (98.33%) as benign and 1 case (1.67%) as malignant. As per **Table - 4, 5**, various studies [5, 6, 7, 8, 9, 10] showed comparable results.

In present study, 32 cases were diagnosed malignant at cytology. Based on this patients were underwent surgery and histopathology confirmed 32 cases (100%) as malignant. As per **Table - 6**, various studies [7, 8, 9, 10] showed comparable results.

Conclusion

The present study was undertaken to study the cytomorphology of various breast lesions and to correlate it with histopathology wherever a surgical excision was done and to establish FNAC as an accurate and reliable preoperative diagnostic tool for breast lesions.

The study consisted of 100 cases (110 aspirates) in which 95 female and 5 were male. Majority of the patients were between 21-40 year age group.

Table - 4: Comparison of various age group incidences of benign breast lesions with other studies.

Age (Years)	Present study	Dr. Venu Anand, et al. (2017) [5]	Neena Chauhan, et al. (2012) [6]
<= 20	17 (28.33%)	23 (16.31%)	12 (14.63%)
21-40	33 (55%)	93 (65.95%)	55 (67.07%)
41-60	8 (13.33%)	23 (16.31%)	13 (15.85%)
61-80	2 (3.33%)	02 (1.41%)	02 (2.43%)
Total	60	141	82

Table – 5: Comparison of cytological benign lesion and histopathology with other studies.

Studies	No. of cytological benign lesion (Total)	Histopathological diagnosis	
		Benign	Malignant
Tiwari M [7]	16	15 (93.75%)	01 (6.25%)
O’Neil S, et al. [8]	166	153 (92.17%)	13(7.83%)
Zhang Qin, et al. [9]	215	213 (99.07%)	02 (0.93%)
A.Z. Mohammed, et al. [10]	61	58 (95.08%)	03 (4.92%)
Present study	60	59 (98.33%)	01 (1.67%)

Table - 6: Comparison of cytological malignant lesion and histopathology with other studies.

Studies	No. of cytological malignant lesion (Total)	Histopathological diagnosis	
		Malignant	Benign
Tiwari M [7]	05	05 (100%)	00 (0.00%)
O’Neil S, et al. [8]	401	398 (99.25%)	03(0.75%)
Zhang Qin, et al. [9]	73	73(100%)	00 (0.00%)
A.Z. Mohammed, et al. [10]	27	27 (100%)	00 (0.00%)
Present study	32	32 (100%)	00 (0.00%)

All the patients presented with a mass in the breast. The following categorizations of FNAC results were observed:

C1-1 cases, C2-60 cases, C3-4, cases C4-3 case, C5-32 cases

The C2 category of 60 benign cases consisted of 20 (majority) cases of fibroadenoma, 3 case of phyllodes tumor, 15 cases of benign proliferative breast disease, 1 case of fibrocystic breast disease, 7 cases of inflammatory lesions and 3 case of intraductal papilloma. Out of 5 cases of gynecomastia, 4 cases were benign and 1 case was malignant (intracystic papillary carcinoma) which confirmed by histopathology.

The C5 category of 32 malignant lesions consisted of 31 cases of infiltrating duct carcinoma- NOS type and 1 case of infiltrating duct carcinoma with medullary type.

The overall sensitivity in the study was 83.33%, specificity was 100%, positive predictive value in diseases was 100% and the false negative percentage was 16.66%.

The study was 100% sensitive in the diagnosis of malignant lesions.

The higher incidence of malignant breast lesions with aged group 41– 60 years of patients.

Thus FNAC is an effective and valid tool as the first line diagnostic modality in the preoperative diagnosis of both benign and malignant lesions.

References

1. Kocjan G, Bourgain C, Fassina A, et al. The role of breast FNAC in diagnosis & clinical management: a survey of current

- practice. *Cytopathology*, 2008; 19: 271-78.
2. Bibbo M, Wilbur DC. *Comprehensive Cytopathology*, 3rd edition, Philadelphia: Saunders Elsevier, 2008; 713-715 (a); 725-750(b).
 3. Bilous M. Breast core needle biopsy: Issues and controversies. *Modern Pathology*, 2010; 23: 536-545.
 4. Rocha PD, Nadkarni NS, Menezes S. Fine needle aspiration biopsy of breast lesions and histopathological correlation. *Acta Cytol.*, 1997; 41(3):705-712.
 5. Dr Venu Anand Dr S. Selvi, Dr. C. Sofiya, Dr. V. Ramya. A study of aspiration cytology of breast and histopathological correlation. *IOSR Journal of dental and medical science*, 2017; 16(12): 58-61.
 6. Neena Chauhan, V.P. Pathak, Meena Harsh, Sunis Saini. Cytohistopathological correlation in palpable breast lesions: Analysis of 200 cases. *Indian Medical Gazette*, December 2012; 473-478.
 7. Tiwari M. Role of FNAC in diagnosis of breast lumps. *Kathmandu University Medical Journal*, 2007; 5: 215-17.
 8. O'Neil S, Castelli M, Gattuso P, Kluskens L, Madsen K, Aranha G. Fine- needle aspiration of 697 palpable breast lesions with histopathologic correlation. *Surgery*, 1997; 122(4): 824–28.
 9. Zhang Qin, Nie Shigui, Chen Yuhua, Zhou Limei. Fine Needle Aspiration Cytology of Breast Lesions: Analysis of 323 Cases. *The Chinese-German Journal of Clinical Oncology*, 2004; 3(3): 172-74.
 10. Mohammad AZ, Edino ST, Ochicha O, Alhassan SU. Value of fine needle aspiration biopsy in preoperative diagnosis of palpable breast lumps in resource-poor countries: a Nigerian experience. *Annals of African Medicine*, 2005; 4(1): 19-22.
 11. Ram Krishan Sharma, Mandeep Radhawa, Neeraj Bisht, N.S. Neki. Fine Needle Aspiration of palpable breast lesions results obtained with PAP MGG with histopathological correlation-hospital based study. *Int. J. Curr. Res. Med. Sci.*, 2017; 3(7): 167-173.
 12. Choi YD, Choi YH, Lee JH, Nam JH, Juhng SW, Choi C. Analysis of Fine Needle Aspiration Cytology of the Breast. *Acta Cytol.*, 2004; 48(6): 801-806.